

## 参考文献・引用文献

### 1 章

- 青木淳一, 1973. 土壤動物学—分類・生態・環境との関係を中心に—, 北隆館.
- 青木淳一, 1983. 自然の診断役・土ダニ, NHK ブックス.
- Coupland, R. T. *et al.*, 1974. Summary of activities, 1967–1974. *Canadian Committee for IBP, Matador Project Technical Report No. 69*, University of Saskatchewan.
- Fittkau, E. J. and Klinge, H., 1973. On biomass and trophic structure of the central Amazonian rain forest Ecosystem. *Biotropica*, **5**, 2–14.
- Gisin, H., 1947. Es wimmelt im Boden von Unbekanntem. *Prisma*, **2** (5), 144–147.
- Hassall, M. *et al.*, 1987. Effects of terrestrial isopods on the decomposition of wood land leaf litter. *Oecologia*, **72**, 597–604.
- Huhta, V., 2007. The role of soil fauna in ecosystems: A historical review. *Pedobiologia*, **50**, 489–495.
- 金子信博, 2007. 土壤生態学入門—土壤動物の多様性と機能—, 東海大学出版会.
- 金子信博, 2012. バイオマス生産による土壤生態系への影響. *Journal of Life Cycle Assessment, Japan*, **8**, 333–337.
- 吉良達夫, 1976. 陸上生態系—概論—, 生態学講座 2, 共立出版.
- Kitazawa, Y., 1971. Biological regionality of the soil fauna and its function in forest ecosystem types. In *Productivity of Forest Ecosystems: Proceedings Brussels Symposium*(Duvigneaud, P. ed.), pp. 485–498, UNESCO.
- 北沢右三, 1974. 志賀山亜高山帯針葉樹林生態系の構造と機能. IBP 成果シンポジウム, 日本学術会議 IBP 特別委員会, pp. 52–56.
- Kitazawa, Y. (ed.), 1977. *Ecosystem Analysis of the Subalpine Coniferous Forest of the Shigayama IBP Area, Central Japan*, University of Tokyo Press.
- 清川昌一ほか, 2014. 地球全史スーパー年表 (日本地質学会監修), 岩波書店.
- Nielsen, U. N., 2019. *Soil Fauna Assemblages: Global to Local Scales*, Cambridge University Press.
- Petersen, H. and Luxton, M. A., 1982. Comparative analysis of soil fauna populations and their role in decomposition processes. *Oikos*, **39**, 288–388.
- Satchell, J. E., 1971. Feasibility study of an energy budget for Meathop Wood. In *Productivity of Forest Ecosystems: Proceedings Brussels Symposium*(Duvigneaud, P. ed.), pp. 619–630, UNESCO.

- Schaefer, M., 1990. The soil fauna of a beech forest on limestone: Trophic structure and energy budget. *Oecologia*, **82**, 128–136.
- Swift, M. J. *et al.*, 1979. *Decomposition in Terrestrial Ecosystems*, Blackwell Scientific Oxford.
- Usher, M. B. *et al.*, 1982. A review of progress in understanding the organization of communities of soil arthropods. *Pedobiologia*, **23**, 126–144.
- Wallwork, J. A., 1970. *Ecology of Soil Animals*, McGraw-Hill.
- 渡辺弘之, 1967. オカダンゴムシのいろいろな温度条件下での摂食量. 日本生態学会誌, **17**, 134–135.
- 渡辺弘之, 1971. 土壌動物の働き. *Edaphologia*, **1**, 7–91.

## 2 章

- Adl, S. M. *et al.*, 2019. Revisions to the classification, nomenclature, and diversity of eukaryotes. *J. Eukaryot. Microbiol.*, **66** (1), 4–119.
- Anderson, F. E. *et al.*, 2017. Phylogenomic analyses of Crassicitellata support major Northern and Southern Hemisphere clades and a Pangaeian origin for earthworms. *BMC Evolutionary Biology*, **17**, 1–18.
- 青木淳一, 1995. 土壌動物を用いた環境診断. 自然環境への影響予測—結果と調査法マニュアル— (沼田 真編), pp. 197–271, 千葉県環境部環境調査課.
- 青木淳一, 2005. だれでもできるやさしい土壌動物のしらべかた—採集・標本・分類の基礎知識—, 102 pp., 合同出版.
- 青木淳一 (編著), 2015. 日本産土壌動物 (第二版) —分類のための図解検索—, 東海大学出版部.
- Bellinger, P. F. *et al.*, 1996–2020. Checklist of the Collembola of the World (ver. 2020 Aug.). Online at <http://www.collembola.org> (Accessed 24 on Sep. 2020) [一澤 (2020) での引用より].
- Brusca, R. C. *et al.*, 2016. *Invertebrates* (3rd ed.), Sinauer Associates.
- Coleman, D. C. *et al.*, 2018. *Fundamentals of Soil Ecology* (3rd ed.), Academic Press.
- Dunlop, J. A. *et al.*, 2021. A summary list of fossil spiders and their relatives. In *World Spider Catalog*, Natural History Museum Bern.  
<http://wsc.nmbe.ch>, version 22.0 (閲覧 2021 年 5 月 4 日)
- Erséus, C. *et al.*, 2020. Phylogenomic analyses reveal a Palaeozoic radiation and support a freshwater origin for clitellate annelids. *Zoologica Scripta*, **49**, 614–640.
- Fernández, R. *et al.*, 2018. Phylogenomics illuminates the backbone of the Myriapoda Tree of Life and reconciles morphological and molecular phylogenies. *Sci. Rep.*, **8** (1), 1–7.
- Giribet, G. and Edgecombe, G. D., 2020. *The Invertebrate Tree of Life*, Princeton University Press.
- 萩原康夫ほか (編著), 2019. 土の中の美しい生き物たち—超拡大写真で見る不思議な生態—, 朝倉書店.

- 一澤 圭, 2020. トビムシの高次分類体系における近年の動向. *Edaphologia*, **108**, 1–21.
- 一澤 圭ほか, 2015. トビムシ目(粘管目) Collembola. 日本産土壌動物(第二版) —分類のための図解検索— (青木淳一編著), pp. 1093–1482, 東海大学出版部.
- Krantz, G. W. and D. E. Walter (eds.), 2009. *A Manual of Acarology* (3rd ed.), Texas Tech University Press.
- 小林元樹, 2021. 環形動物門の高次系統に関する概説. *Edaphologia*, **109**, 9–17.
- Lamsdell, J. C. *et al.*, 2020. Air breathing in an exceptionally preserved 340-million-year-old sea scorpion. *Curr. Biol.*, **30** (21), 4316–4321.
- Martín-Durán, J. M. *et al.*, 2016. The larval nervous system of the penis worm *Priapulius caudatus* (Ecdysozoa). *Philos. Trans. R. Soc. B*, **371** (1685), 20150050.
- 南谷幸雄, 2014–2022. 日本産ミミズ大図鑑. <https://japanese-mimizu.jimdofree.com/> (閲覧 2022 年 5 月 10 日)
- Misof, B. *et al.*, 2014. Phylogenomics resolves the timing and pattern of insect evolution. *Science*, **346**, 763–767.
- 永野昌博・澤島拓夫(編), 2009. 雪・森・農のめぐみとつながりを考えるシリーズ 1, 森を支える小さな戦士—落ち葉の下の生き物たち—. 十日町市里山科学館「森の学校」キヨロロ, 十日町市.
- Nolan, E. D. *et al.*, 2020. Developmental gene expression as a phylogenetic data class: Support for the monophyly of Arachnopolmonata. *Dev. Genes Evol.*, **2020**, 1–17.
- 大久保憲秀ほか, 2015. ササラダニ亜目. 日本産土壌動物(第二版) —分類のための図解検索— (青木淳一編著), pp. 346–717, 東海大学出版部.
- Regier, J. C. *et al.*, 2010. Arthropod relationships revealed by phylogenomic analysis of nuclear protein-coding sequences. *Nature*, **463** (7284), 1079–1083.
- Schneider, C. *et al.*, 2011. Unexpected diversity in Neelipleona revealed by molecular phylogeny approach (Hexapoda, Collembola). *Soil Organisms*, **83**, 383–398.
- 芝 実, 2015. ケダニ亜目. 日本産土壌動物(第二版) —分類のための図解検索— (青木淳一編著), pp. 201–313, 東海大学出版部.
- 島野智之, 2014. 生物指標の手法と原生生物への応用. 原生生物フロンティア—その生物学と工学— (洲崎敏伸編), pp. 103–114, 化学同人.
- 島野智之, 2018a. 原生生物. 実践土壌学シリーズ 2 土壌生態学 (金子信博編), 朝倉書店.
- 島野智之, 2018b. ダニ類の高次分類体系の改訂について—高次分類群の一部和名改称—. 日本ダニ学会誌, **27**, 51–68.
- 島野智之, 2018c. なぜダニ類はクモガタ類の中で最も種数が多いのか? タクサ, **44**, 4–14.
- 島野智之, 2021. ダニ類の高次分類群とその和名. *Edaphologia*, **109**, 34–35.
- 島野智之・青木淳一\*, 2015. ダニ目. 日本産土壌動物(第二版) —分類のための図解検索— (青木淳一編著), pp. 148–150, 東海大学出版部. \*著者順は正誤表で訂正.
- 島野智之・高久 元(編著), 2016. ダニのはなし—人間との関わり—, 朝倉書店.
- Subías, L. S., 2021. Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes, Oribatida) del mundo (except fósils). [Original publishing in 2004. *Graellista*, **60**, 3–305.] <http://bba.bioucm.es/cont/docs/RO.1.pdf> (閲覧 2021 年 4 月 8 日)
- Weigert, A. *et al.*, 2014. Illuminating the base of the annelid tree using transcriptomics.

- Mol. Biol. Evol.*, **31**, 1391–1401. doi:10.1093/molbev/msu080
- Woese, C. R. *et al.*, 1990. Towards a natural system of organisms: Proposal for the domains Archaea, Bacteria, and Eucarya. *PNAS*, **87**, 4576–4579.
- Xiong, Y. *et al.*, 2008. Molecular phylogeny of Collembola inferred from ribosomal RNA genes. *Mol. Phylogenetics Evol.*, **49**, 728–735.
- 矢崎裕規・島野智之, 2020. 真核生物の高次分類体系の改訂—Adl *et al.* (2019) について—タクサ, **48**, 71–83.
- Zhang, F. and Deharveng, L., 2015. First instar tibiotarsal chaetotaxy supports the Entomobryidae and Symphypleona (Collembola) forming a cluster in a phylogenetic tree. *Zootaxa*, **3955**, 487–504.
- Zhang, Z.-Q., 2013. Phylum Arthropoda. *Zootaxa*, **3703**, 17–26.

### 3 章

- Anderson, J. M. and Bignell, D. E., 1980. Bacteria in the food, gut contents and faeces of the litter-feeding millipede *Glomeris marginata* (Villers). *Soil Biol. Biochem.*, **12**, 251–254.
- Anderson, J. *et al.*, 1983. Nitrogen and cation mobilization by soil fauna feeding on leaf litter and soil organic matter from deciduous woodlands. *Soil Biol. Biochem.*, **15**, 463–467.
- Barois, I. *et al.*, 1998. Andosol-forming process linked with soil fauna under the perennial grass *Mulhembergia macroura*. *Geoderma*, **86**, 241–260.
- Berg, M. *et al.*, 2001. Community food web, decomposition and nitrogen mineralisation in a stratified Scots pine forest soil. *Oikos*, **94**, 130–142.
- Bonkowski, M. *et al.*, 2000. Microbial-faunal interactions in the rhizosphere and effects on plant growth. *Eur. J. Soil Biol.*, **36**, 135–147.
- Bonkowski, M., 2009. Rhizosphere fauna: The functional and structural diversity of intimate interactions of soil fauna with plant roots. *Plant and Soil*, **321**, 213–233.
- Bradford, M. A. *et al.*, 2002. Impacts of soil faunal community composition on model grassland ecosystems. *Science*, **298**, 615–618.
- Breznak, J. A. and Brune, A., 1994. Role of Microorganisms in the digestion of lignocellulose by termites. *Ann. Rev. Entomol.*, **39**, 453–487.
- Brown, V. K. and Gange, A. C., 1992. Secondary plant succession: How is it modified by insect herbivory? *Vegetatio*, **101**, 3–13.
- Brown, G. G. *et al.*, 2000. Regulation of soil organic matter dynamics and microbial activity in the drilosphere and the role of interactions with other edaphic functional domains. *Eur. J. Soil Biol.*, **36**, 177–198.
- Cárcamo, H. A. *et al.*, 2000. Influence of millipedes on litter decomposition, N mineralization, and microbial communities in a coastal forest in British Columbia, Canada. *Can. J. For. Res.*, **30**, 817–826.
- Crowther, T. W. and A’Bear, A. D., 2012. Impacts of grazing soil fauna on decomposer

- fungi are species-specific and density-dependent. *Fung. Ecol.*, **5**, 277–281.
- Crowther, T. W. *et al.*, 2011. Outcomes of fungal interactions are determined by soil invertebrate grazers. *Ecol. Lett.*, **14**, 1134–1142.
- de Graaff, M. A. *et al.*, 2015. A meta-analysis of soil biodiversity impacts on the carbon cycle. *Soil*, **1**, 257–271.
- Faber, J. H. and Verhoef, H. A., 1991. Functional differences between closely-related soil Arthropods with respect to decomposition processes in the presence or absence of Pine tree roots. *Soil Biol. Biochem.*, **23**, 15–23.
- Forey, E. *et al.*, 2011. Importance of earthworm–seed interactions for the composition and structure of plant communities: A review. *Acta Oecol.*, **37**, 594–603.
- Frouz, J. *et al.*, 2008. Effects of the exotic Crustacean, *Armadillidium vulgare* (Isopoda), and other macrofauna on organic matter dynamics in soil microcosms in a hardwood forest in a central Florida. *Florida Entomol.*, **91**, 328–331.
- Griffiths, B. S., 1990. A comparison of microbial-feeding nematodes and protozoa in the rhizosphere of different plants. *Biol. Fertil. Soils*, **9**, 83–88.
- Hanlon, R. and Anderson, J., 1979. The effects of Collembola grazing on microbial activity in decomposing leaf litter. *Oecologia*, **38**, 93–99.
- Hättenschwiler, S. and Gasser, P., 2005. Soil animals alter plant litter diversity effects on decomposition. *PNAS*, **102**, 1519–1524.
- Hector, A. *et al.*, 2000. Consequences of the reduction of plant diversity for litter decomposition: Effects through litter quality and microenvironment. *Oikos*, **90**, 357–371.
- Heemsbergen, D. A. *et al.*, 2004. Biodiversity effects on soil processes explained by interspecific functional dissimilarity. *Science*, **306**, 1019–1020.
- 本郷裕一・大熊盛也, 2008. シロアリ腸内共生微生物群の多様性とゲノム解析. *J. Env. Biotech.*, **8**, 29–34.
- Hooper, D. U. *et al.*, 2005. Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecol. Mono.*, **75**, 3–35.
- Ineson, P. *et al.*, 1982. Effects of collembolan grazing upon nitrogen and cation leaching from decomposing leaf litter. *Soil Biol. Biochem.*, **14**, 601–605.
- Jones, C. G. *et al.*, 1994. Organisms as ecosystem engineers. *Oikos*, **69**, 373–386.
- 金子信博, 2007. 土壤生態学入門, 東海大学出版会.
- Kautz, G. *et al.*, 2002. Does *Porcellio scaber* (Isopoda: Oniscidea) gain from coprophagy? *Soil Biol. Biochem.*, **34**, 1253–1259.
- King, A. J. *et al.*, 2010. Molecular insight into lignocellulose digestion by a marine isopod in the absence of gut microbes. *PNAS*, **107**, 5345–5350.
- Lattaud, C. *et al.*, 1998. The diversity of digestive systems in tropical geophagous earthworms. *Appl. Soil Ecol.*, **9**, 189–195.
- Lavelle, P., 1997. Faunal activities and soil processes: Adaptive strategies that determine ecosystem function. *Adv. Ecol. Res.*, **27**, 91–132.
- Lavelle, P. *et al.*, 2004. Effects of earthworms on soil organic matter and nutrient dynamics at a landscape scale over decades. *Earthworm Ecology* (2nd ed.), pp.

- 145–160, CRC Press.
- Lavelle, P. and Martin, A. V., 1992. Small-scale and large-scale effects of endogeic earthworms on soil organic matter dynamics in soil and the humid tropics. *Soil Biol. Biochem.*, **24**, 1491–1498.
- Lavelle, P. and Spain, A. V., 2001. *Soil Ecology*, Kluwer Academic Publishers.
- Liiri, M. *et al.*, 2002. Soil processes are not influenced by the functional complexity of soil decomposer food webs under disturbance. *Soil Biol. Biochem.*, **34**, 1009–1020.
- Liu, T. *et al.*, 2019. Earthworms coordinate soil biota to improve multiple ecosystem functions. *Curr. Biol.*, **29**, 3420–3429.
- Maaß, S. *et al.*, 2015. Functional role of microarthropods in soil aggregation. *Pedobiologia*, **58**, 59–63.
- Milcu, A. *et al.*, 2006. Earthworms (*Lumbricus terrestris*) affect plant seedling recruitment and microhabitat heterogeneity. *Funct. Ecol.*, **20**, 261–268.
- Moore, J. C. *et al.*, 2003. Top-down is bottom-up: Does predation in the rhizosphere regulate aboveground dynamics? *Ecology*, **84**, 846–857.
- Newell, K., 1984a. Interaction between two decomposer Basidiomycetes and a collembolan under Sitka spruce: Grazing and its potential effects on fungal distribution and litter decomposition. *Soil. Biol. Biochem.*, **16**, 235–239.
- Newell, K., 1984b. Interaction between two decomposer Basidiomycetes and a collembolan under Sitka spruce: Distribution, abundance and selective grazing. *Soil. Biol. Biochem.*, **16**, 227–233.
- Nielsen, U. N., 2019. *Soil Fauna Assemblages: Global to Local Scales*, Cambridge University Press.
- Nielsen, U. N. *et al.*, 2011. Soil biodiversity and carbon cycling: A review and synthesis of studies examining diversity-function relationships. *Eur. J. Soil Sci.*, **62**, 105–116.
- 新島溪子, 1984. キシヤヤスデの大発生. 森林立地, **26**, 25–32.
- Schultz, J. C. *et al.*, 2013. Flexible resource allocation during plant defense responses. *Front. Plant Sci.*, **4**, 324.
- Setälä, H. *et al.*, 1990. Effects of soil fauna on leaching of nitrogen and phosphorus from experimental systems simulating coniferous forest floor. *Biol. Fertil. Soils*, **10**, 170–177.
- 島野智之, 2018. 第2章 原生生物. 実践土壌学シリーズ2 土壌生態学 (金子信博編), pp. 14–30, 朝倉書店.
- Soler, R. *et al.*, 2007. Impact of foliar herbivory on the development of a root-feeding insect and its parasitoid. *Oecologia*, **152**, 257–264.
- Sugimoto, A. *et al.*, 2000. Global impact of termites on the carbon cycle and atmospheric trace gases. In *Termites: Evolution, Sociality, Symbioses, Ecology* (Abe, T. *et al.*, eds.), pp. 409–435, Kluwer Academic Publishers.
- Tilman, J. *et al.*, 1997. The influence of functional diversity and composition on ecosystem processes. *Science*, **277**, 1300–1302.
- 角田智詞, 2018. 第8章 植物の根系と植食昆虫の関係. 実践土壌学シリーズ2 土壌生態学

- (金子信博編), pp. 101–115, 朝倉書店.
- Wagg, C. *et al.*, 2014. Soil biodiversity and soil community composition determine ecosystem multifunctionality. *PNAS*, **111**, 5266–5270.
- Wardle, D. A. *et al.*, 2004. Ecological linkages between aboveground and belowground biota. *Science*, **304**, 1629–1633.
- Watanabe, H. and Tokuda, G., 2010. Cellulolytic systems in insects. *Ann. Rev. Entomol.*, **55**, 609–632.
- Wurst, S. *et al.*, 2008. Intraspecific variation in plant defense alters effects of root herbivores on leaf chemistry and aboveground herbivore damage. *J. Chem. Ecol.*, **34**, 1360–1367.
- Zimmer, M., 2002. Nutrition in terrestrial isopods (Isopoda: Oniscidea): An evolutionary ecological approach. *Biol. Rev.*, **77**, 455–493.
- Zimmer, M. *et al.*, 2005. Do woodlice and earthworms interact synergistically in leaf litter decomposition? *Funct. Ecol.*, **19**, 7–16.

#### 4 章

- Kobayashi, M. *et al.*, 2017. Spore dissemination by mycophagous adult drosophilids. *Ecol. Res.*, **32** (4), 621–626.
- 大園享司, 2018. 基礎から学べる菌類生態学, 共立出版.
- 相良直彦, 1989. きのこと動物—ひとつの地下生物学—, 築地書館. (2021年に新訂版刊行)
- Sawahata, T. *et al.*, 2000. Number and food habit of springtails on wild mushrooms of three species of Agaricales. *Edaphologia*, (66), 21–33.
- 津田 格, 2012. キノコと昆虫を利用する線虫たち—森をめぐるミクロな世界—. 微生物生態学への招待 (二井一禎ほか編), pp. 127–145, 京都大学学術出版会.

#### 5.1 節

- 石塚小太郎・皆越ようせい, 2014. ミミズ図鑑, 全国農村教育協会.
- 南谷幸雄, 2014–2022. 日本産ミミズ大図鑑. <https://japanese-mimizu.jimdofree.com/> (閲覧 2022年5月10日)
- 中村好男, 2015. ミミズ綱 (貧毛綱). 日本産土壌動物 (第二版) —分類のための図解検索— (青木淳一編著), pp. 1893–1902, 東海大学出版会.

#### 5.2 節

- Angst, G. *et al.*, 2019. Earthworms act as biochemical reactors to convert labile plant compounds into stabilized soil microbial necromass. *Communications Biology*, **2**, 441.

- FAO *et al.*, 2020. State of knowledge of soil biodiversity—Status, challenges, and potentialities. Report 2020, FAO.
- Kaneda, S. *et al.*, 2016. Soil temperature and moisture-based estimation of rates of soil aggregate formation by the endogeic earthworm *Eisenia japonica* (Michaelsen, 1892). *Biol. Fertil. Soils*, **52**, 789–797.
- 金子信博, 2018. 保全農業と土壤動物. 実践土壌学シリーズ 2 土壤生態学 (金子信博編), pp. 147–161, 朝倉書店.
- Kawaguchi, T. *et al.*, 2011. Mineral nitrogen dynamics in the casts of epigeic earthworms (*Metaphire hilgendorfi*: Megascolecidae). *Soil Sci. Plant Nutr.*, **57**, 387–395.
- Lavelle, P. *et al.*, 1997. Soil function in a changing world: The role of invertebrate ecosystem engineers. *Eur. J. Soil Sci.*, **33**, 159–193.
- Lee, K. E. and Foster, R. C., 1991. Soil fauna and soil structure. *Aust. J. Soil Res.*, **29**, 745–775.
- Oades, J. M., 1993. The role of biology in the formation, stabilization and degradation of soil structure. *Geoderma*, **56**, 377–400.

### 5.3 節

- Butt, K., 2011. The earthworm inoculation unit technique: Development and use in soil improvement over two decades. In *Biology of Earthworms* (Soil Biology) (Karaca, A. ed.), pp. 87–105, Springer.
- Curl, E. A. and Lartey, R. T., 1996. Role of soil fauna in biological control of *Rhizoctonia*. In *Rhizoctonia Species: Taxonomy, Molecular Biology, Ecology, Pathology and Disease Control* (Sneh, B. *et al.* eds.), pp. 495–506, Kluwer Academic Publishers.
- Edwards, C. A. and Bohlen, P. J., 1996. *Biology and Ecology of Earthworms* (3rd ed.), Chapman and Hall.
- van Groenigen, J. W. *et al.*, 2014. Earthworms increase plant production: A meta-analysis. *Sci. Rep.*, **4**, 6365.
- 伊藤豊彰ほか, 2011. 冬期湛水・有機栽培水田の土壤動物：イトミミズの生態と機能 (シンポジウム 有機農業を巡る土壤微生物・小動物). *土と微生物*, **65**, 94–99.
- Kikuchi, E. *et al.*, 1975. Surveys of the fauna and flora in the water and soil of paddy fields. *Rep. Inst. Agric. Res. Tohoku University*, **26**, 25–35.
- Kikuchi, E. and Kurihara, Y., 1977. In vitro studies on the effects of tubificids on the biological, chemical and physical characteristics of submerged ricefield soil and overlying water. *Oikos*, **29**, 348–356.
- Klironomos, J. N. *et al.*, 1999. Reproductive significance of feeding on saprobic and arbuscular mycorrhizal fungi by the collembolan, *Folsomia candida*. *Func. Ecol.*, **13**, 756–761.
- 小荒井 晃, 2004. 水田における一年生広葉雑草の発生生態とイネ品種によるそれらへの抑制



- 効果に関する研究. 中央農研研究報告, **5**, 59–102.
- 栗原 康, 1983. イトミミズと雑草—水田生態系解析への試み (1) —. 化学と生物, **21**, 243–249.
- 栗原 康・菊地永祐, 1983. イトミミズと雑草 II イトミミズの波及効果. 化学と生物, **21**, 324–327.
- Lee, K. E., 1985. *Earthworms their Ecology and Relationships with Soils and Land Use*, 411 pp., Academic Press.
- Mitra, P. and Kaneko, N., 2017. Impact of aquatic earthworms on methane emission reduction from the paddy field soil in Japan. *J. Agri. Sci.*, **9**, 36–46.
- 中村好男ほか, 1991. 福島県から採集された作物病原糸状菌を摂食する中型土壤動物. *Edaphologia*, **45**, 49–54.
- 白石啓義ほか, 1993. トビムシ移入によるアズキ幼苗の白紋羽病抑制効果. 東北農業研究センター研究報告, **46**, 139–140.
- 白石啓義ほか, 2005. 育苗箱へのトビムシ導入によるアブラナ科野菜の苗立枯れ症抑制. 東北農業研究センター研究報告, **103**, 53–61.
- 白石啓義・中村好男, 1994. 乾性土壤動物による土壤病害抑制効果—トビムシ *Folsomia hidakana* と糸状菌由来の土壤病害—. 東北農業研究, **47**, 171–172.
- Tajbakhsh, J. *et al.*, 2011. Vermicompost as a biological soil amendment. In *Biology of Earthworms* (Soil Biology) (Karaca, A. ed.), pp. 215–228, Springer.

#### 5.4 節

- Ballabio, C. *et al.*, 2018. Copper distribution in European topsoils: An assessment based on LUCAS soil survey. *Sci. Total Environ.*, **636**, 282–298.
- Briones, M. J. I. and Schmid, O., 2017. Conventional tillage decreases the abundance and biomass of earthworms and alters their community structure in a global meta-analysis. *Glob. Change Biol.*, **23**, 4396–4419.
- Hendrix, P. F. and Edwards, C. A., 2004. Earthworms in agroecosystems: Research approaches. In *Earthworm Ecology* (Edwards, C. A. ed.), pp. 287–295, CRC PRESS.
- 久馬一剛, 1984. 新土壌学, 271 pp., 朝倉書店.
- Lee, K. E., 1985. *Earthworms their Ecology and Relationships with Soils and Land Use*, Academic Press.
- 望月武雄ほか, 1975. 農薬の散布によって重金属類の蓄積したリング園土壌の動物生態学的研究 (第 1 報): 青森県津軽地方のリング園土壌中の銅, 鉛, ヒ素含量と大形土壤動物相について. 日本土壌肥科学雑誌, **46**, 45–50.
- 中村好男, 1998. ミミズと土と有機農業—「地球の虫」のはたらき—, 創森社.
- Naveed, M. *et al.*, 2014. Simultaneous loss of soil biodiversity and functions along a copper contamination gradient: When soil goes to sleep. *Soil Sci. Soc. Am. J.*, **78**, 1239–1250.
- Paoletti, M. G., 1999. The role of earthworms for assessment of sustainability and as bioindicators. *Agri. Ecosys. Environ.*, **74**, 137–155.

- Pelosi, C. *et al.*, 2014. Pesticides and earthworms. A review. *Agron. Sustain. Dev.*, **34**, 199–228.
- Wardle, D. A., 1995. Impacts of disturbance on detritus food webs in agro-ecosystems of contrasting tillage and weed management practices. *Adv. Ecol. Res.*, **26**, 105–185.

## 6 章

- 工藤慎一ほか, 2014. ヤスデ 2 種で新たに確認された抱卵行動. 昆虫 (ニューシリーズ), **17**, 135–137.
- Marek, P. E. *et al.*, 2021. The first true millipede—1306 legs long. *Scientific Reports*, **11**, 23126.
- 三好保徳, 1951. 雄が抱卵するヒラタヤスデ. 採集と飼育, **13**, 54–55.
- 村上好央, 1965. 日本産普通多足類の後胚発生 XVIII イヨハガヤスデの生活史 (2). 動物学雑誌, **74**, 31–37.
- Nijijima, K. *et al.*, 2021. Eight-year periodical outbreaks of the train millipede. *Royal Society Open Science*, **8** (1), 201399.
- 篠原圭三郎, 1974. 多足類の採集と観察, グリーンブックス 12, pp. 2–84, ニューサイエンス社.
- 高桑良興, 1954. 日本産倍足類総説, pp. 1–17, 日本学術振興会.
- 田辺 力, 2001. 多足類読本, pp. 96–161, 東海大学出版会.
- Toyota, A. *et al.*, 2006. Soil ecosystem engineering by the train millipede *Parafontaria laminata* in a Japanese larch forest. *Soil Biol. Biochem.*, **38** (7), 1840–1850.
- Toyota, A. and Kaneko, N., 2012. Plant productivity is temporally enhanced by soil fauna depending on the life stage and abundance of animals. In *Forest Ecosystems: More than Just Trees* (Blanco, J. and Lo, Y.-H. eds.), 11, pp. 253–264, InteckOpen.

## 7 章

- Guisan, A. *et al.*, 2017. *Habitat Suitability and Distribution Models: With Applications in R*, Cambridge University Press.
- Hatano, M. *et al.*, 2015. Ontogenetic metamorphosis and extreme sexual dimorphism in lumpsuckers: *Eumicrotremus asperrimus*, *Cyclopteropsis bergi* and *Cyclopteropsis lindbergi*, may be synonymous. *J. Fish. Biol.*, **86**, 1121–1128.
- 岩崎貴也ほか, 2016. 分子系統地理学に生態ニッチモデリングがもたらす新展開と課題. 植物地理・分類研究, **64**, 1–15.
- Jackson, M. *et al.*, 2017. Using nested PCR to improve detection of earthworm eDNA in Canada. *Soil Biol. Biochem.*, **113**, 215–218.
- Karasawa, S., 2016. Eleven nominal species of *Burmoniscus* are junior synonyms of *B.*

- kathmandius* (Schmalfuss, 1983)(Crustacea, Isopoda, Oniscidea). *Zookeys*, **607**, 1–24.
- 唐沢重考ほか, 2021. アマミサソリモドキ *Typopeltis stimpsonii* (Wood, 1862) の分布状況および系統地理学的考察. *Edaphologia*, **109**, 19–31.
- Karasawa, S. and Honda, M., 2016. Taxonomic study of the *Burmoniscus ocellatus* complex (Crustacea, Isopoda, Oniscidea) in Japan shows genetic diversification in the Southern Ryukyus, Southwestern Japan. *Zool. Sci.*, **29**, 527–537.
- 木村政昭, 2002. 琉球弧の成立と生物の渡来, 沖縄タイムス社.
- Lomolino, M. V. *et al.*, 2017. *Biogeography—Biological Diversity across Space and Time* (5th ed.), Sinauer Associates.
- 曾田貞滋, 2013. 新オサムシ学—生態から進化まで—, 北隆館.

## 8 章

- Beltran-Garcia, M. J. *et al.*, 1997. Volatile compounds secreted by the oyster mushroom (*Pleurotus ostreatus*) and their antibacterial activities. *J. Agric. Food Chem.*, **45**, 4049–4052.
- Bengtsson, G. *et al.*, 1991. Selective odor perception in the soil collembola *Onychiurus armatus*. *J. Chem. Ecol.*, **17**, 2113–2125.
- Berendsen, R. L. *et al.*, 2013. Effects of the mushroom-volatile 1-octen-3-ol on dry bubble disease. *Appl. Microbiol. Biotechnol.*, **97**, 5535–5543.
- Bitzer, C. *et al.*, 2004. Benzoic acid derivatives in a hypogastrurid collembolan: Temperature-dependent formation and biological significance as deterrents. *J. Chem. Ecol.*, **30**, 1591–1602.
- Brückner, A. *et al.*, 2018. Track the snack olfactory cues shape foraging behaviour of decomposing soil mites (Oribatida). *Pedobiologia*, **66**, 74–80.
- Chitarra, G. S. *et al.*, 2004. Germination of *Penicillium paneum* conidia is regulated by 1-octen-3-ol, a volatile self-inhibitor. *Appl. Environ. Microbiol.*, **70** (5), 2823–2829.
- Davis, T. S. *et al.*, 2013. Microbial volatile emissions as insect semiochemicals. *J. Chem. Ecol.*, **39**, 840–859.
- Dettner, K. *et al.*, 1996. Chemical defense of giant springtail *Tetrodotoxiphora bielaensis* (Waga) (Insecta: Collembola). *J. Chem. Ecol.*, **22**, 1051–1074.
- Ditengou, F. A. *et al.*, 2015. Volatile signalling by sesquiterpenes from ectomycorrhizal fungi reprogrammes root architecture. *Nat. Commun.*, **6**, 6279.
- Fialho, M. B. *et al.*, 2012. Nematicidal effect of volatile organic compounds (VOCs) on the plant-parasitic nematode *Meloidogyne javanica*. *Summa. Phytopathol.*, **38**, 152–154.
- Freire, E. S. *et al.*, 2012. Volatile substances produced by *Fusarium oxysporum* from coffee rhizosphere and other microbes affect *Meloidogyne incognita* and *Arthrobotrys conoides*. *J. Nematol.*, **44**, 321–328.
- Frey-Klett, P. *et al.*, 2011. Bacterial-fungal interactions: Hyphens between agricultural,

- clinical, environmental, and food microbiologists. *Microbiol. Mol. Biol. Rev.*, **75**, 583–609.
- 藤井義晴・濱野満子, 2003. アレロパシー：植物-植食者-天敵相互作用系における植物情報化学物質の機能. *日生氣誌*, **40**, 49–54.
- Grimme, E. *et al.*, 2007. Comparison of *Muscodor albus* volatiles with a biorational mixture for control of seedling diseases of sugar beet and root-knot nematode on tomato. *Plant Dis.*, **91**, 220–225.
- Hågvar, S. *et al.*, 1995. Long distance, directional migration on snow in a forest collembolan, *Hypogastrura socialis* (Uzel). *Acta. Zool. Fennica.*, **196**, 200–205.
- Hågvar, S. *et al.*, 2000. Navigation and behavior of four Collembola species migrating on the snow surface. *Pedobiologia*, **44**, 221–233.
- Hung, R. *et al.*, 2013. *Arabidopsis thaliana* as a model system for testing the effect of *Trichoderma volatile* organic compounds. *Fungal. Ecol.*, **6**, 19–26.
- Kalisz, S. *et al.*, 2021. Allelopathy in pervasive in invasive plants. *Biol. Invasions*, **23**, 367–371.
- Kaneda, S. and Kaneko, N., 2004. The feeding preference of a collembolan (*Folsomia candida* Willem) on ectomycorrhiza (*Pisolithus tinctorius* (Pers.)) varies with mycelial growth condition and vitality. *Appl. Soil Ecol.*, **27**, 1–5.
- Kline, D. L., 1994. Olfactory attractants for mosquito surveillance and control: 1-octen-3-ol. *J. Amer. Mosq. Control Assoc.*, **10**, 280–287.
- Klironomos, J. N. and Hart, M. M., 2001. Food-web dynamics: Animal nitrogen swap for plant carbon. *Nature*, **410**, 651–652.
- Leinaas, H. P., 1981. Cyclomorphosis in *Hypogastrura lapponica* (Axelson, 1902) (= *H. frigida* (Axelson, 1905) syn. nov.) (Collembola, Poduridae). Morphological adaptations and selection for winter dispersal. *Zeitschrift für Zoologische Systematik und Evolutionsforschung*, **19**, 278–285.
- Meents, A. K., 2019. Volatile DMNT systemically induces jasmonate-independent direct anti-herbivore defense in leaves of sweet potato (*Ipomoea batatas*) plants. *Sci. Rep.*, **9**, 17431. doi:10.1038/s41598-019-53946-0
- Messer, C. *et al.*, 2000. Chemical deterrents in podurid Collembola. *Pedobiologia*, **44**, 210–220.
- Nagaki, M. *et al.*, 2011. Antibacterial and antifungal activities of isoprenoids. *Trans. Mat. Res. Soc. Jpn*, **36**, 55–58.
- 中森泰三, 2009. 菌食性トビムシの餌選択と菌類の防御. *日菌報*, **50**, 71–78.
- 岡崎久美子, 2019. きこの由来揮発性物質による作物保護技術の開発—きこの香りで作物を病気から守る—. *日本農業学会誌*, **44**, 69–70.
- Okull, D. O. *et al.*, 2003. Antifungal activity of 10-oxo-trans-8-decenoic acid and 1-octen-3-ol against *Penicillium expansum* in potato dextrose agar medium. *J. Food Prot.*, **66**, 1503–1505.
- Porco, D. and Deharveng, L., 2007. 1, 3-Dimethoxybenzene, a chemotaxonomic marker for the Neanurinae subfamily (Collembola). *Biochem. Syst. Ecol.*, **35**, 160–161.
- Salmon, S. *et al.*, 2019. Chemical communication in springtails: A review of facts and

- perspectives. *Biol. Fertil. Soils*, **55**, 425–438.
- Sawahata, T., 2008. *Tricholoma matsutake* 1-octen-3-ol and methyl cinnamate repel mycophagous *Proisotoma minuta* (collembola: insecta). *Mycorrhiza*, **18**, 111–114.
- 塩尻かおりほか, 2002. 植物–植食者–天敵相互作用系における植物情報化学物質の機能. 応動昆, **46**, 117–133.
- Strobel, G., 2006. *Muscodor albus* and its biological promise. *J. Ind. Microbiol. Biotechnol.*, **33**, 514–522.
- Sun, X. *et al.*, 2015. Effect of volatiles versus exudates released by germinating spores of *Gigaspora margarita* on lateral root formation. *Plant Physiol. Biochem.*, **97**, 1–10.
- Wener, S. *et al.*, 2016. Belowground communication: Impacts of volatile organic compounds (VOCs) from soil fungi on other soil-inhabiting organisms. *Appl. Microbiol. Biotechnol.*, **100**, 8651–8665.
- Wood, W. F. *et al.*, 2001. 1-octen-3-ol, a banana slug antifeedant from mushroom. *Biochem. Syst. Ecol.*, **29**, 531–533.
- Zettel, J. and Zettel, U., 2008. Manche mögen's kalt: die Biologie des "Schneeflohs" *Ceratophysella sigillata* (Uzel, 1891), einer winteraktiven Springschwanzart (Collembola: Hypogastruridae). In *Mitteilungen der Naturforschenden Gesellschaft in Bern*, Neue Folge, Band 65, pp. 79–110, Naturforschende Gesellschaft in Bern.

## 9 章

- Aoki, J. and Ohnishi, J., 1974. New species and record of oribatid mites from Hokkaido, North Japan. *Bulletin of the National Science Museum, Tokyo*, **17** (2), 149–156.
- Blum, M. S., 1996. Semiochemical parsimony in the Arthropoda. *Annu. Rev. Entomol.*, **41**, 353–374.
- Brückner, A., *et al.*, 2017. Storage and release of hydrogen cyanide in a chelicerate (*Oribatula tibialis*). *PNAS*, **114**, 3469–3472.
- Carrel, J. and Eisner, T., 1984. Spider sedation induced by defensive chemicals of millipede prey. *PNAS*, **81**, 806–810.
- Cloudsley-Thompson, J. L., 1949. The enemies of myriapods. *Naturalist*, **831**, 137–141.
- Dadashipour, M. *et al.*, 2015. Discovery and molecular and biocatalytic properties of hydroxynitrile lyase from an invasive millipede, *Chamberlinius hualienensis*. *PNAS*, **112**, 10605–10610.
- Deml, R. and Huth, A., 2000. Benzoquinones and hydroquinones in defensive secretions of tropical millipedes. *The Science of Nature*, **87**, 80–82.
- Duffey, S. and Blum, M., 1977. Phenol and guaiacol biosynthesis, detoxication and function in a polydesmid millipede *Oxidus gracilis*. *Insect Biochem.*, **7**, 57–66.
- Eisner, H. *et al.*, 1967. Defense mechanisms of Arthropods. XX. Quantitative assessment of hydrogen cyanide production in two species of millipedes. *Psyche*, **74**,

- 107–117.
- Eisner, T. *et al.*, 1978. Defensive secretions of millipedes. In *Arthropod Venoms* (Bettini, S. ed.), Handbook of Experimental Pharmacology, pp. 41–72, Springer-Verlag.
- Eisner, T. *et al.*, 2005. *Secret Weapons: Defenses of Insects, Spiders, Scorpions and Other Many-legged Creatures*, Belknap Press.
- Heethoff, M. and Rall, B. C., 2015. Reducible defence: Chemical protection alters the dynamics of predator–prey interactions. *Chemoecology*, **25**, 53–61.
- Herbert, D., 2000. Dining on diplopods: Remarkable feeding behavior in chlamydephorid slugs (Mollusca: Gastropoda). *J. Zool.*, **251**, 1–5.
- Jałoszyński, P. and Olszanowski, Z., 2016. Feeding of two species of Scydmaeninae “hole scrapers”, *Cephennium majus* and *C. ruthenum* (Coleoptera: Staphylinidae), on oribatid mites. *Eur. J. Entomol.*, **113**, 372–386.
- Kuwahara, Y. *et al.*, 1975. Alarm pheromone of the cheese mite, *Tyrophagus putrescentiae* (Schrank) (Acarina, Acaridae), *Experientia*, **31**, 1115–1116.
- 桑原保正, 1999. ヤスデの化学防御. 環境昆虫学—行動・生理・化学生態— (日高敏隆・松本義明監修, 本田計一ほか編), pp. 291–298, 東京大学出版会.
- Kuwahara, Y. *et al.*, 2002. 2-Nitroethenylbenzenes as natural products in millipede defense secretions. *Naturwissenschaften*, **89**, 308–310.
- Kuwahara, Y., 2004. Chemical ecology of astigmatid mites. In *Advances in Insect Chemical Ecology* (Cardè, R. T. and Millar, J. G. eds.), pp. 76–109, Cambridge University Press.
- Kuwahara, Y. *et al.*, 2007. Detection of a neotropical frog alkaloid spiropyrrolizidine 236 from a Japanese polyzoniid millipede *Kiusiozonium okai* as a major defense component together with polyzonimine and nitropolyzonimine. *Jpn. J. Environ. Ent. Zool.*, **18**, 91–96.
- Kuwahara, Y. *et al.*, 2011. Release of hydrogen cyanide via a post-secretion Schotten-Baumann reaction in defensive fluids of polydesmoid millipedes. *J. Chem. Ecol.*, **37**, 232–238.
- Kuwahara, Y. *et al.*, 2017. Hydrogen peroxide as a new defensive compound in “benzoyl cyanide” producing polydesmid millipedes. *Sci. Nat.*, **104**, 19.
- Maruno, G. *et al.*, 2012. Chemical ecology of astigmatid mites LXXXVII. *S*-(+)-isopiperitenone: Re-identification of the alarm pheromone as the female sex pheromone in *Tyrophagus similis* (Acari: Acaridae). *J. Chem. Ecol.*, **38**, 36–41.
- Meinwald, J. *et al.*, 1975. Nitropolyzonamine: A spirocyclic compound from the defensive glands of a millipede (*Polyzonium rosalbum*). *Tetrahedron Lett.*, **28**, 2367–2370.
- Mitaka, Y. *et al.*, 2017. Multi-functional roles of a soldier-specific volatile as a worker arrestant, primer pheromone and an antimicrobial agent in a termite. *Proceedings of the Royal Society B*, **284**, 20171134.
- Nishimura, K. *et al.*, 2002. Chemical ecology of astigmatid mites. LXIV. The alarm pheromone neral functions as an attractant in *Schwiebea elongata* (Banks) (Acari: Acaridae). *Appl. Entomol. Zool.*, **37**, 13–18.
- Rasputnig, G., 2006. Chemical alarm and defence in the oribatid mite *Collohmanna*

- gigantea* (Acari: Oribatida). *Exp. Appl. Acarol.*, **39**, 177–194.
- Saporito, R. *et al.*, 2003. A siphonotid millipede (*Rhinotus*) as the source of spiropyrrrolizidine oximes of dendrobatid frogs. *J. Chem. Ecol.*, **29**, 2781–2786.
- Saporito, R. A. *et al.*, 2007. Oribatid mites as a major dietary source for alkaloids in poison frogs. *PNAS*, **104**, 8885–8890.
- Schildknechte, H. and Wenneis, W., 1967a. Über Arthropoden-Abwehrstoffe. XX. Structuraufklärung des Glomerins. *Z. Natur.*, **21b**, 552–556.
- Schildknechte, H. and Wenneis, W., 1967b. Über Arthropoden-Abwehrstoffe. XXV. Anthranils aure als Precursor der Arthropoden-alkaloide Glomerin und Homoglomerin. *Tetrahedron Lett.*, **8**, 1815–1818.
- Shear, W. *et al.*, 2011. Glomerin and homoglomerin from the North American pill millipede *Onomeris sinuata* (Loomis, 1943) (Diplopoda, pentazonia, glomeridae). *Int. J. Myriapodology*, **4**, 1–10.
- Shear, W., 2015. The chemical defenses of millipedes (Diplopoda) : Biochemistry, physiology and ecology. *Biochem. Syst. Ecol.*, **61**, 78–117.
- Shimano, S. *et al.*, 2002. Geranial: The alarm pheromone in the nymphal stage of the oribatid mite, *Nothrus palustris*. *J. Chem. Ecol.*, **28**, 1831–1837.
- Shimizu, N. *et al.*, 2001. Aggregation pheromone activity of the female sex pheromone,  $\beta$ -Acaridial, in *Caloglyphus polyphyllae* (Acari: Acaridae). *Biosci. Biotechnol. Biochem.*, **65**, 1724–1728.
- Shimizu, N. *et al.*, 2012. *n*-Hexyl laurate and fourteen related fatty-acid esters: New secretory compounds from the julid millipede, *Anaulaciulus* sp. *J. Chem. Ecol.*, **38**, 23–28.
- Shimizu, N. *et al.*, 2012. The absolute configuration of chrysomelidial: A widely distributed defensive component among Oribotritiid mites (Acari: Oribatida). *J. Chem. Ecol.*, **38**, 29–35, 2012.
- 清水伸泰, 2018. ダニと昆虫の多機能フェロモンとその活用術. 昆虫と自然: 特集「昆虫のフェロモン」, **53**, 4–7.
- Smolanoff, J. *et al.*, 1975. Polyzonimine: A novel terpenoid insect repellent produced by a millipede. *Science*, **188**, 734–736.
- Takada, W. *et al.*, 2005. Schelorbitid mites as the source of pumiliotoxins in dendrobatid frogs. *J. Chem. Ecol.*, **31**, 2403–2415.
- Tatjana, L. *et al.*, 2014. Quinones and non-quinones from the defensive secretion of *Uniciger transsilvanicus* (Verhoeff, 1899) (Diplopoda, Julida, Julidae), from Serbia. *Arch. Biol. Sci.*, **66**, 385–390.
- Vujisić, L. V. *et al.*, 2013. Chemistry of the sternal gland secretion of the Mediterranean centipede *Himantarium gabrielis* (Linnaeus, 1767) (Chilopoda: Geophilomorpha: Himantariidae). *Naturwissenschaften*, **100**, 861–870.
- Wood, W. *et al.*, 2000. Buzonamine, a new alkaloid from the defensive secretion of the millipede, *Buzonium crassipes*. *Biochem. Syst. Ecol.*, **28**, 305–312.
- Wyatt, T. D., 2014. *Pheromones and Animal Behavior: Chemical Signals and Signatures* (2nd ed.), Cambridge University Press.

Yamaguchi, T. *et al.*, 2017. A novel cytochrome P450, CYP3201B1, is involved in (*R*)-mandelonitrile biosynthesis in a cyanogenic millipede. *FEBS Open Bio*, **7**, 335–347.

## 10 章

青木淳一（編著），2015. 日本産土壌動物（第二版）—分類のための図解検索—，東海大学出版会.

Bokhorst, S. *et al.*, 2012. Extreme winter warming events more negatively impact small rather than large soil fauna: Shift in community composition explained by traits not taxa. *Glob. Change Biol.*, **18**, 1152–1162. doi:10.1111/j.1365-2486.2011.02565.x

Hishi, T. *et al.*, 2007. The feeding habits of collembola along decomposition gradients using stable carbon and nitrogen isotope analyses. *Soil Biol. Biochem.*, **39**, 1820–1823. doi:10.1016/j.soilbio.2007.01.028

菱 拓雄ほか，2010. 九州大学北海道演習林の天然落葉広葉樹林およびカラマツ人工林における斜面方位に着目した土壌と大型土壌動物の特徴. 九大演報, **91**, 1–6.

Hishi, T. *et al.*, 2019. Taxonomy, distribution and trait data sets of Japanese Collembola. *Ecol. Res.*, **34**, 444–445. doi:10.1111/1440-1703.12022

菱 拓雄，2022. 形質生態学によるトビムシ群集構造の解明. *Edaphologia*, 印刷中.

Hishi, T. *et al.*, 2022. Topography is more important than forest type as a determinant for functional trait composition of Collembola community. *Pedobiologia*, **90**, 150776. doi:10.1016/j.pedobi.2021.150776

Hubbell, S. P., 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*, Princeton University Press. (平尾聡秀ほか（訳），2009. 群集生態学—生物多様性学と生物地理学の統一中立理論—，文一総合出版.)

Hyodo, F. *et al.*, 2006. Estimation of the longevity of C in terrestrial detrital food webs using radiocarbon ( $^{14}\text{C}$ ): How old are diets in termites? *Funct. Ecol.*, **20**, 385–393. doi:10.1111/j.1365-2435.2006.01081.x

Hyodo, F. *et al.*, 2015. Dependence of diverse consumers on detritus in a tropical rain forest food web as revealed by radiocarbon analysis. *Funct. Ecol.*, **29**, 423–429. doi:10.1111/1365-2435.12357

Janssens, F. and Lawrence, N., 2002–2012. Are Collembola terrestrial Crustacea? <http://www.collembola.org/publicat/crustacn.htm> (閲覧 2022 年 6 月 13 日)

McGill, B. J. *et al.*, 2006. Rebuilding community ecology from functional traits. *Trends Ecol. Evol.*, **21**, 178–185. doi:10.1016/j.tree.2006.02.002

Okuzaki, Y. *et al.*, 2009. Vertical heterogeneity of a forest floor invertebrate food web as indicated by stable-isotope analysis. *Ecol. Res.*, **24**, 1351–1359. doi:10.1007/s11284-009-0619-0

Petersen, H., 2002. General aspects of collembola ecology at the turn of millennium. *Pedobiologia*, **46**, 246–260.



- Vellend, M., 2016. *The Theory of Ecological Communities*, Princeton University Press.  
(松岡俊将ほか (訳), 2019. 群集生態学の理論: 4 つのルールで読み解く生物多様性, 共立出版.)
- Wardle, D. A. *et al.*, 2004. Ecological linkages between aboveground and belowground biota. *Science*, **304**, 1629–1633.
- Widenfalk, L. *et al.*, 2016. Small-scale Collembola community composition in a pine forest soil—Overdispersion in functional traits indicates the importance of species interactions. *Soil Biol. Biochem.*, **103**, 52–62. doi:10.1016/j.soilbio.2016.08.006

## 11 章

- 青木淳一, 1973. 土壤動物学—分類・生態・環境との関係を中心に—, 北隆館.
- 青木淳一, 1995. 土壤動物を用いた環境診断. 自然環境への影響予測—結果と調査法マニュアル— (沼田 真編), pp. 197–271, 千葉県環境部環境調査課.
- 青木淳一, 2005. だれでもできるやさしい土壤動物のしらべかた—採集・標本・分類の基礎知識—, 102 pp., 合同出版.
- 萩原康夫, 2019. 第 4 章 土壤動物を対象とした自然観察会の案内. 土の中の美しい生き物たち—超拡大写真で見る不思議な生態— (萩原康夫ほか編著), pp.139–152, 朝倉書店.
- 金子信博, 2007. 土壤生態学入門—土壤動物の多様性と機能—, 東海大学出版会.
- 加藤良一ほか, 2013. 使い捨てカイロとペットボトルを用いた簡易型ツルグレン装置. 山形大学紀要, **15**, 341–352.
- ミュージアムパーク茨城県自然博物館 (編), 2007. 第 39 回企画展「ありんこアントの大冒険—土の中の生きものを探せ!—」, 34 pp.
- 永野昌博・澤島拓夫 (編), 2009. 雪・森・農のめぐみとつながりを考えるシリーズ 1. 森を支える小さな戦士—落ち葉の下の生き物たち—. 十日町市里山科学館「森の学校」キョロロ, 十日町市.
- 島野智之, 2019. 第 1 章 土壤動物とは何か. 土の中の美しい生き物たち—超拡大写真で見る不思議な生態— (萩原康夫ほか編著), pp.1–10, 朝倉書店.

## コラム 4

- Ikeda, H. *et al.*, 2020. A comparison of latitudinal species diversity patterns between riverine and terrestrial earthworms from the North American temperate zone. *Journal of Biogeography*, **47**, 1373–1382.
- Reynolds, J. W., 2008. Sparganophilidae (Annelida, Oligochaeta) distribution in North America and elsewhere, revisited. *Megadrilologica*, **12**, 125–143.
- Reynolds, J. W. and Wetzell, M. J., 2008. Terrestrial Oligochaeta (Annelida: Clitellata) in North America, including Mexico, Puerto Rico, Hawaii, and Bermuda. *Megadrilologica*, **12**, 157–210.
- Rota, E. *et al.*, 2014. First time in Italy. Is the elusive aquatic megadrile *Sparganophilus*

Benham 1892 (Annelida, Clitellata) accelerating its dispersal in Europe? *Journal of Limnology*, **73**, 482–489.

Rota, E. *et al.*, 2016. Mitochondrial evidence supports a Nearctic origin for the spreading limicolous earthworm *Sparganophilus tamesis* Benham, 1892 (Clitellata, Sparganophilidae). *Contributions to Zoology*, **85**, 113–119.